

Population frequencies of transposable elements in selfing and outcrossing *Caenorhabditis* nematodes

ELIE S. DOLGIN^{1*}, BRIAN CHARLESWORTH¹ AND ASHER D. CUTTER^{2*}

¹ Institute of Evolutionary Biology, School of Biological Sciences, University of Edinburgh, King's Buildings, Edinburgh, UK

² Department of Ecology and Evolutionary Biology, University of Toronto, Toronto, ON, Canada

(Received 23 April 2008 and in revised form 2 July 2008)

Summary

Population genetics theory predicts that differences in breeding systems should be an important factor in the dynamics of selfish genetic elements, because of different intensities of selection on both hosts and elements. We examined population frequencies of transposable elements (TEs) in natural populations of the self-fertilizing nematode *Caenorhabditis elegans* and its outcrossing relative *Caenorhabditis remanei*. We identified a Tc1-like class of elements in the *C. remanei* genome with homology to the terminal inverted repeats of the *C. elegans* Tc1 transposon, which we name mTc1. We measured levels of insertion polymorphism for all 32 Tc1 elements present in the genome sequence of the *C. elegans* N2 strain, and 16 mTc1 elements from the genome sequence of the *C. remanei* PB4641 strain. We show that transposons are less polymorphic and segregate at higher frequencies in *C. elegans* compared with *C. remanei*. Estimates of the intensity of selection based on the population frequencies of polymorphic elements suggest that transposons are selectively neutral in *C. elegans*, but subject to purifying selection in *C. remanei*. These results are consistent with a reduced efficacy of natural selection against TEs in selfing populations, but may in part be explained by non-equilibrium TE dynamics.

1. Introduction

The selfish DNA hypothesis proposes that the abundance of transposable elements (TEs) in natural populations of their hosts is the consequence of a balance between the proliferation of elements by transposition and natural selection acting to remove insertions from the population (Doolittle & Sapienza, 1980; Orgel & Crick, 1980). Accordingly, any factor that modulates the strength of selection or rates of transposition should have an important effect on the distribution and dynamics of element insertions. At the genomic level, the impact of differences in the efficacy of selection on the patterning of TEs can be seen in the genomes of many organisms, and elements have been

found to accumulate differentially among chromosomal regions in association with recombination rate differences (Charlesworth & Langley, 1989; Duret *et al.*, 2000; Boissinot *et al.*, 2001; Bartolomé *et al.*, 2002; Rizzon *et al.*, 2002, 2003), base composition patterns (Lander *et al.*, 2001) and gene density (Medstrand *et al.*, 2002).

At the population level, a potentially important factor for TE dynamics is the host breeding system (Hickey, 1982; Charlesworth & Charlesworth, 1995; Wright & Schoen, 1999; Morgan, 2001). When populations do not mate randomly, elements can be affected in a number of ways with different expected consequences. Self-fertilization can lead to element accumulation, because inbreeding reduces the effective population size experienced by the host, and smaller populations tend to accumulate more elements (Charlesworth & Charlesworth, 1983; Brookfield & Budge, 1997; Wright & Schoen, 1999; Morgan, 2001). In an inbreeding population, insertions will also tend to become homozygous: If the deleterious effects of TEs are caused by ectopic recombination between

* Corresponding authors: Elie S. Dolgin, Institute of Evolutionary Biology, School of Biological Sciences, University of Edinburgh, King's Buildings, Edinburgh EH9 3JT, UK. Tel: +44 131 650 5476. Fax: +44 131 650 6564. e-mail: elie.dolgin@ed.ac.uk; and Asher D. Cutter, Department of Ecology and Evolutionary Biology, University of Toronto, 25 Harbord Street, Toronto, ON, M5S 3G5, Canada. Tel: +1 416 978 4602. Fax: +1 416 978 5878. e-mail: asher.cutter@utoronto.ca

dispersed, heterozygous insertions (Montgomery *et al.*, 1991), selection may be more effective in out-crossing populations. But selfing can also lead to the containment of TEs, and if individual insertions are harmful because they disrupt genes (Finnegan, 1992), inbreeding should enhance the efficacy of natural selection against TEs because of higher homozygosity (Charlesworth & Charlesworth, 1995). In addition, the complete stochastic loss of elements in small, self-fertilizing populations is also possible (Wright & Schoen, 1999; Morgan, 2001). A final effect of selfing is that lower levels of genetic exchange in inbred hosts might influence the evolution of transposition rates themselves, leading to conditions favouring TEs that reduce their own activity levels (Charlesworth & Langley, 1986; Charlesworth & Charlesworth, 1995).

Comparisons of the population frequencies of TEs in natural populations have been performed within and between many animal species, including fruitflies (Charlesworth & Langley, 1989; Petrov *et al.*, 2003; Bartolomé & Maside, 2004; Franchini *et al.*, 2004), mosquitoes (O'Brochta *et al.*, 2006; Boulesteix *et al.*, 2007), midges (Zampicini *et al.*, 2004), several fish species (Takasaki *et al.*, 1997; Duvernell & Turner, 1999; Neafsey *et al.*, 2004) and humans (Batzer & Deininger, 2002; Bennett *et al.*, 2004). These studies have shed light on the selective constraints experienced by TEs, but, to date, the only empirical investigations of the effects of breeding systems on TE dynamics have focused on closely related self- and cross-pollinating plant species (Wright *et al.*, 2001; Tam *et al.*, 2007). Both of these studies found a significant effect of the breeding system – insertion polymorphism levels were lower and individual elements segregated at higher frequencies in self-fertilizing species. Whether these conclusions apply to TEs in selfing populations more generally remains to be seen.

In the present study, we perform the first systematic survey of TE frequencies in two species of the nematode genus *Caenorhabditis* – the model organism *Caenorhabditis elegans*, which reproduces primarily as a self-fertilizing hermaphrodite, and the related out-crossing species, *Caenorhabditis remanei* – and test the role of breeding systems in driving TE dynamics. TEs make up about 12% of the *C. elegans* genome, with the most active and best characterized being type II DNA transposons of the Tc1/*mariner* superfamily (see the review by Bessereau, 2006). These elements, which are among the most widespread DNA transposons (Plasterk *et al.*, 1999), were named after the superfamily's two best-studied members: Tc1, the first TE identified in *C. elegans* (Emmons *et al.*, 1983; Liao *et al.*, 1983), and *mariner*, which was first described in *Drosophila mauritiana* (Jacobson *et al.*, 1986). These transposons have highly diverged primary sequences, but all Tc1/*mariner* elements probably derive from a common ancestor, and they share many common

features including flanking terminal inverted repeats (TIRs) and similar modes of transposition (Plasterk *et al.*, 1999). The genome of the canonical *C. elegans* strain N2 contains 32 copies of Tc1 (Fischer *et al.*, 2003), but the copy number is strain-specific, and different wild strains harbour different numbers of Tc1 transposons, ranging from around 30 copies in most strains to upwards of 300 copies in others (Emmons *et al.*, 1983; Liao *et al.*, 1983; Egilmez *et al.*, 1995; Hodgkin & Doniach, 1997). Although these elements do not show germline transposition in N2, they can be activated in the germline by mutation of a single gene in the N2 strain (Collins *et al.*, 1987; Ketting *et al.*, 1999), activity can arise spontaneously in a non-mutator strain (Babity *et al.*, 1990), and some natural isolates show high rates of germline activity (Emmons *et al.*, 1983; Eide & Anderson, 1985).

The *C. remanei* genome is much more poorly characterized than that of *C. elegans* and, as such, much less is known about its TEs. Nonetheless, sequences homologous to Tc1 transposons are widely distributed among nematodes (Moerman & Waterston, 1989), polymorphic Tc1-like elements have been found segregating in other nematode species (Hoekstra *et al.*, 1999), and probes designed for the *C. elegans* Tc1, Tc2, and Tc3 elements hybridize weakly in *C. remanei* (Abad *et al.*, 1991). In addition, the associated species *Caenorhabditis briggsae*, which is more closely related to *C. remanei* than *C. elegans*, contains transposons similar to Tc1, and these show polymorphic hybridization patterns (Harris *et al.*, 1990). Here, we describe a class of Tc1-like transposons in *C. remanei*, and examine both *C. elegans* and *C. remanei* for transposon population frequencies – the degree to which individual transposons segregate in natural populations. We show that levels of insertion polymorphism significantly differ between the two species and evaluate the possible role of mating system in driving TE dynamics.

2. Materials and methods

(i) *Transposable elements*

The canonical N2 strain of *C. elegans* contains 32 copies of the Tc1 DNA transposon (Fischer *et al.*, 2003). We quantified the population frequencies for each of these Tc1 elements in a global sample of *C. elegans* strains. Transposon locations were taken from the genomic positions identified by Fischer *et al.* (2003). For *C. remanei*, we identified a homologous class of transposons by using RepeatMasker (<http://www.repeatmasker.org>) and the Tc1 and *C. briggsae* Tc1 sequences as the basis for Blastn queries of the *C. remanei* genome draft 15.0.1 for strain PB4641, which was sequenced using plasmid and fosmid libraries to a depth of 9.2× and assembled using the

Table 1. *C. elegans* and *C. remanei* strains used in the present study

Strains	Location (nearest city)	Source
<i>C. elegans</i>		
N2	Bristol, UK	CGC
AB1, AB4	Adelaide, Australia	CGC
KR314	Vancouver, Canada	CGC
PX174	Portland, OR, USA	CGC
PS2025, CB4854	Altadena, CA, USA	CGC
CB4855	Palo Alto, CA, USA	CGC
CB4857	Claremont, CA, USA	CGC
TR403	Madison, WI, USA	CGC
CB4856	Hawaii, USA	CGC
JU1088	Kakegawa, Japan	MAF
JU1171	Concepcion, Chile	MAF
LKC34	Madagascar	MAF
ED3040	Johannesburg, South Africa	ED
ED3046, ED3052	Ceres, South Africa	ED
ED3057	Limuru, Kenya	ED
ED3077	Nairobi, Kenya	ED
JU258	Ribeiro Frio, Madeira, Portugal	MAF
JU776, JU780, JU799	Lisbon, Portugal	MAF
RC301	Freiburg, Germany	CGC
MY1, MY2, MY6, MY12	Münster, Germany	CGC
RW7000	Bergerac, France	CGC
JU314, JU323	Merlet, France	MAF
JU365	Franconville, France	MAF
JU393, JU394, JU395, JU398, JU400, JU402, JU407, JU439	Hermanville, France	MAF
<i>C. remanei</i>		
PB4641	Brooklyn, NY, USA	CGC
SB146	Freiburg, Germany	CGC
PB207, PB210, PB211, PB213, PB215, PB219, PB242, PB243, PB244, PB247, PB249, PB252, PB253, PB275	Dayton, OH, USA	SB

CGC, *Caenorhabditis* Genetics Center; MAF, Marie-Anne Félix; ED, Elie Dolgin; SB, Scott Baird.

PCAP whole-genome assembly program (Washington University Genome Sequencing Center). The transposons we describe are similar to Tc1 and Tcb1 in the terminal regions, but they are smaller and may not be autonomous, so we have called them mTc1 elements, where ‘m’ stands for miniature (N. Jiang, personal communication) and ‘re’ is an abbreviation for *C. remanei*. We determined the consensus sequence for mTc1 transposons using BioEdit, and measured the population frequencies for a random subset of ‘full-length’ elements.

(ii) *Nematode populations and molecular methods*

For *C. elegans*, we tested for the presence or absence of all 32 of the Tc1 elements found in the N2 strain for each of 39 *C. elegans* strains, defined as isohermaphrodite lines from which we obtain single haplotypes (Table 1). Thirty-one strains were chosen to provide worldwide representation of geographic locations where this species has been found, and an additional eight strains were selected from a single sampling locality in Hermanville, France. Genetic diversity levels

at nuclear loci in Hermanville samples are similar to those found on a global level (Barrière & Félix, 2005; Cutter, 2006). For *C. remanei*, we quantified the presence or absence for 16 elements identified in PB4641 for the inbred strain SB146 and for 14 isofemale strains from Wright State University Biological Preserve in Dayton, OH, USA (Table 1). We selected the strains from a single geographic locality for *C. remanei* because this population of *C. remanei* does not show deviation from demographic equilibrium, and so this sample is likely to be representative of the species (Cutter *et al.*, 2006; Cutter, 2008).

For *C. elegans*, genomic DNA was isolated from whole plates of worms with the Puregene DNA Purification kit (D-7000A, Gentra Systems). For *C. remanei*, we performed whole-genome amplification of single males using the REPLI-g Mini kit (Qiagen). Diluted aliquots of the DNA samples were then used as templates for standard PCR reactions. We used DNA from the sequenced strains of *C. elegans* (N2) and *C. remanei* (PB4641) as positive controls in our PCR amplification of Tc1 and mTc1 elements, respectively.

(iii) *Population frequency assays*

We determined the presence or absence of individual transposon insertions in all strains using pairs of PCR primers in the flanking regions on either side of the transposon insertion sites, coupled with a third primer matching an internal portion of the transposon sequence (e.g. Bartolomé & Maside, 2004). The PCRs performed with flanking primers produce unique sized bands that show whether the transposon is present or absent, and the reactions with the internal primers serve as additional tests to confirm transposon presence. For *C. elegans*, a single internal primer was used for a conserved region in 31 of the 32 Tc1 insertion sites, while a different internal primer was used for clone C50H2, due to a 701-bp deletion in the transposon (Fischer *et al.*, 2003). For *C. remanei*, PCR failure rates were higher, presumably due to greater levels of nucleotide polymorphism, so we used two sets of flanking primers and unique internal primers for most transposon insertion sites. We used different combinations of primers until every strain was successfully amplified. Primer sets were designed for 21 mTc1 insertions in PB4641, but for five of these, only the positive control (PB4641) worked for any primer combination. Consequently, we restrict our analysis to the 16 mTc1 insertion sites that yielded interpretable data. The primer sequences are provided in supplementary Table S1.

Due to high levels of somatic transposition (Emmons & Yesner, 1984), we could not differentiate homozygotes and heterozygotes when an insertion was detected. As a result, transposon presence functioned as a dominant marker. For *C. elegans*, we calculated the population frequencies by assuming that all transposons are homozygous, owing to the high degree of selfing. Therefore, the frequency of each transposon is simply equal to the number of strains for which the insertion was detected, divided by the total number of strains (excluding N2). For *C. remanei*, we first discriminated between X-linked *vs.* autosomal insertions by inferring the chromosomal location of each insertion using Blastn of the unique flanking regions against the 2005 preliminary assembly of the *C. remanei* genome in Wormbase (<http://www.wormbase.org>). Based on wobble-aware bulk aligner (WABA) alignments of the resulting *C. remanei* contigs to the *C. elegans* genome, we then inferred the likely syntenic chromosome. Because we used single male DNA preparations for *C. remanei*, heterozygosity of X-linked insertions is not possible, so the frequency of such mTc1 insertions is simply the observed population frequency. For autosomal insertions, we assumed random mating, which seems to be appropriate for the Ohio sample (Cutter *et al.*, 2006; Cutter, 2008), and calculated the frequency of each transposon according to Hardy–Weinberg expectations as

one minus the square root of the frequency of strains that lacked the insertion. Frequency calculations were restricted to the 14 Ohio samples, with the inbred SB146 strain excluded from the analysis.

(iv) *Estimating the strength of selection*

We used the diffusion approximation methods derived by Petrov *et al.* (2003) to estimate the probability that an insertion is at a particular population frequency, and calculated a maximum likelihood estimate of the selection coefficient acting on the transposons. We calculated 95% confidence intervals around the maximum likelihood scores to obtain measures of the intensity of natural selection, $N_e s$, assuming semi-dominance (i.e. $h=0.5$), effective population sizes of 10^4 and 10^6 for *C. elegans* and *C. remanei*, respectively (Cutter, 2006, 2008; Cutter *et al.*, 2006), and that all transposons have independent effects and are subject to the same strength of selection, s . Qualitative conclusions were unaffected by increasing or decreasing the effective population sizes one order of magnitude (results not shown). This analysis only considers segregating elements (i.e. not fixed) and, by using posterior probability functions, accounts for the fact that, by only studying insertions present in the reference sequenced strain, we have pre-sampled transposons in proportion to their population frequencies (Petrov *et al.*, 2003). For *C. elegans*, the analysis was done for the entire collection of 39 strains, for the 8 strains from Hermanville on their own, and for 30 random ‘scattered sample’ subsets with a single strain selected from each geographic location (23 strains in total). This subsampling approach of taking a single strain from each locality may approximate a homogeneously mixing population for a large number of localities connected by migration (Wakeley, 2003; Cutter, 2006; Matsen & Wakeley, 2006). For *C. remanei*, the analysis was limited to the Ohio population.

This analysis assumes a large number of independently segregating insertions at transposition–selection balance. To test this, we measured the level of neutral nucleotide diversity among Tc1 transposons within the genome of the canonical N2 strain and total nucleotide diversity among ‘full-length’ mTc1 transposons within the PB4641 *C. remanei* genome using DnaSP 4.10.9 (Rozas *et al.*, 2003). At copy number equilibrium, population genetic theory predicts that the level of neutral nucleotide diversity among active transposons is equal to $4N_e\Lambda\mu$, where μ is the neutral mutation rate, Λ is the average number of active transposons per haploid genome and $2N_e\Lambda$ is the effective population size of the transposon family (Brookfield, 1986; Sánchez-García *et al.*, 2005). Therefore, silent site diversity between transposons is expected to equal the average haploid copy number




Fig. 1. (A) Consensus nucleotide sequence of the mTcre1 *C. remanei* transposon, with the TIRs shown in boxes. (B) Schematic representation of mTcre1 showing homology to Tc1, Tcb1 and Tc7. Arrows represent the TIRs, with the shaded areas corresponding to regions of sequence similarity.

multiplied by the average genomic silent-site diversity, assuming that the mutation rate in transposons is similar to that of the genome as a whole (Brookfield, 1986). We also measured levels of linkage disequilibrium among transposons for the 39 *C. elegans* strains (excluding N2) and for the 14 Ohio *C. remanei* strains, using the squared correlation between pairs of sites (r^2). Because our sample size, n , for each species is small, the effective population size multiplied by the recombination rate is expected to be much greater than the sample size, and the expected linkage disequilibrium, $E(r^2) \approx 1/n$ (Weir & Hill, 1980). Therefore, we subtract $1/n$ from our r^2 -values, and compare deviations from expectations for the two species.

With the well-annotated genome of *C. elegans*, we also tested for correlations between the population frequency of Tc1 transposons and various aspects of the genomic environment: local recombination rate, gene density, transposon polymorphism levels [single nucleotide polymorphism (SNPs) and indels], and whether the transposon is inserted in an intron. Recombination rates based on the nearest ten loci were taken from Cutter & Payseur (2003), the numbers of SNPs and indels were obtained from Fischer *et al.* (2003), and gene density estimates and whether the transposon was found in an intron were determined using Wormbase (<http://www.wormbase.org>).

3. Results

(i) *C. remanei* transposons

We identified a class of transposons, which we refer to as mTcre1 transposons, exhibiting high sequence similarity to the ends of the TIRs of Tc1 and Tcb1 elements. Figure 1 shows the consensus mTcre1 sequence, which matches 17 of the outer 18 bp in the TIR of Tc1, and 30 of the outer 31 bp in the Tcb1 sequence. The mTcre1 elements have longer TIRs and a shorter total length than either Tc1 or Tcb1. The unique 86 bp internal sequence is probably too short to contain an open reading frame; however, the conserved portion of the TIRs probably contains a transposase binding site, because the Tc1 transposase binding site was identified within the outer portion of the Tc1 inverted repeats (Vos & Plasterk, 1994). In this way, mTcre1 elements might more closely resemble other Tc-family transposons such as *C. elegans* Tc7 elements, which are shorter than Tc1 and rely on Tc1-derived transposase activity (Rezsohazy *et al.*, 1997). Furthermore, mTcre1 elements are flanked by TA dinucleotides, just like Tc1 and Tcb1, which presumably result from target site sequence duplication upon integration (van Luenen *et al.*, 1994). We identified 81 mTcre1 transposons in total, inferred from similarity to the TIRs, although this is probably an underestimate of the true number of mTcre1




Fig. 2. For legend see opposite page.

elements in the PB4641 genome. Of these, 80% are 'full-length' (i.e. similar in length to the consensus mTcrl sequence), 11% contain large (>100 bp) deletions and 9% have complex insertions making them at least 400 bp larger.

(ii) Transposon population frequencies

Whether each transposon was present or absent in all the strains tested is shown in Fig. 2. Individual *C. elegans* strains were significantly more likely than *C. remanei* strains to harbour the transposons tested (Mann–Whitney $U=568.5$, $P<10^{-6}$); on average, ~80% of the 32 Tc1 transposons found in N2 were present in any given wild strain of *C. elegans*, while *C. remanei* strains only had around half of the mTcrl elements detected in PB4641. The population frequencies of transposons were also much higher for Tc1 elements in *C. elegans* than for mTcrl elements in *C. remanei* (Mann–Whitney $U=454.5$, $P<10^{-6}$), as the frequency spectrum was skewed towards high-frequency elements in *C. elegans* (Fig. 3). A greater proportion of insertions was fixed for Tc1 (9 of 32 elements) than for mTcrl (1 of 16), although this difference was not statistically significant (Fisher's Exact Test, $P=0.13$). For *C. elegans*, the population frequency of Tc1 transposons did not correlate with recombination rate, gene density, polymorphism levels or whether the transposon was found in an intron (all Spearman's $|\rho|<0.20$, $P>0.30$). The presence/absence status of transposons also showed no strong signature of geographic structuring in *C. elegans*, consistent with other studies of genetic diversity (Denver *et al.*, 2003; Barrière & Félix, 2005; Haber *et al.*, 2005; Cutter, 2006; Dolgin *et al.*, 2008).

(iii) Strength of selection

Using the observed population frequency distributions of polymorphic transposons to estimate the maximum likelihood strength of selection, we found evidence for purifying selection ($N_{e}s<0$) in *C. remanei* but not in *C. elegans*. For mTcrl elements in *C. remanei*, $N_{e}s=-2.2$ (95% confidence interval: $-3.9\leq N_{e}s\leq -0.6$). In contrast, when considering all *C. elegans* strains tested for Tc1 elements, $N_{e}s=5.9$ ($1.3\leq N_{e}s\leq 27.3$), suggesting that Tc1 elements are subject to positive selection in *C. elegans*. This seems




Fig. 3. Histogram of population frequencies of *C. elegans* Tc1 transposons (light shaded bars) and *C. remanei* mTcrl transposons (dark shaded bars).

unlikely, however, because Tc1 transposons cause large mutagenic effects when transpositionally active in the germline (Plasterk & van Luenen, 1997; Bégin & Schoen, 2006, 2007).

The positive $N_{e}s$ value for *C. elegans* arises from the large number of high-frequency transposons (see Fig. 3), which could be an artefact of population structure if many strains shared a recent N2-like common ancestor. To attempt to remove the effects of population subdivision, we created random subsets of 'scattered samples' with only a single strain from each geographic locality (Wakeley, 2003; Cutter, 2006; Matsen & Wakeley, 2006). This analysis gave an average strength of selection, $N_{e}s\approx 4$. However, heterogeneity between strains from the same location led to 95% confidence intervals for different scattered random subsets to imply no significant difference from neutrality in 7 of 30 subsets, and positive selection in the remaining 23. Overall, the range of lower 95% confidence intervals for the 30 random scattered subsamples spanned the range -0.2 to 1.1 . Considering just the eight strains from a single locality in Hermanville, France, we found no significant difference from neutrality: $N_{e}s=1.5$ ($-1.1\leq N_{e}s\leq 16.8$). Further, the lack of evidence for purifying selection in *C. elegans* is not an artefact of the assumption of complete homozygosity. If we assume Hardy–Weinberg

Fig. 2. Transposon frequencies for *C. elegans* (A) and *C. remanei* (B). Dark squares indicate presence and white squares indicate absence of the transposon. Population frequencies of each transposon calculated for the 39 *C. elegans* strains (excluding N2) and for the 14 Ohio *C. remanei* strains are shown along the bottom; the fraction of elements found for each strain is shown along the right; and the position of each transposon, denoted by the clone (*C. elegans*) or supercontig (*C. remanei*) and the linkage group, is shown along the top. For *C. remanei* supercontig 951, the WABA alignment was inconclusive between linkage groups IV and V.

ratios of allele frequencies, as we did for *C. remanei*, we still observe no significant departure from neutral expectations when considering all 39 strains: $N_{es}=0.3$ ($-1.1 \leq N_{es} \leq 2.8$).

The maximum likelihood method for calculating the strength of selection assumes that transposons are at copy number equilibrium. To evaluate this assumption, we contrasted predictions of nucleotide diversity and linkage disequilibrium with observed values (see the 'Materials and Methods' section). For *C. remanei*, total nucleotide diversity among 'full-length' mTcrl elements in the PB4641 genome was $\pi=18.2\%$ and $\theta=21.3\%$. These diversity measures include parts of the transposon that might experience selection, including the inverted repeats and conserved region, so silent site diversity is presumably even greater. Nonetheless, these values are much greater than the 3.6–4.7% overall silent-site diversity for *C. remanei* (Cutter *et al.*, 2006; Cutter, 2008), consistent with equilibrium expectations (Brookfield, 1986). In contrast, we found that nucleotide diversity at silent sites among the 32 *C. elegans* Tc1 elements in the N2 genome was $\pi_{si}=0.19\%$ and $\theta_{si}=0.76\%$. These values are similar to the overall silent-site diversity of 0.2–0.3% (Cutter, 2006), and not $\sim 30\times$ greater as predicted by equilibrium theory (Brookfield, 1986). Furthermore, this analysis may be conservative as extra rounds of DNA replication during transposition could enhance the nucleotide mutation rate of TEs above that of the rest of the host's genome. We also found Tajima's $D=-1.89$ ($P<0.05$), indicating an excess of low-frequency variants among Tc1 elements, and the mean deviation of linkage disequilibrium from expectation was greater for *C. elegans*: 0.056 between all pairs of sites and 0.112 for intrachromosomal comparisons, compared with r^2 -deviations for *C. remanei* of 0.025 and 0.010, respectively. A greater proportion of r^2 -values was also significant by Fisher's exact tests in *C. elegans* than *C. remanei* (Fisher's exact test between species, $P=0.002$). Together, these measures all suggest that the assumption of copy number equilibrium might be violated for *C. elegans*.

4. Discussion

Phylogenetic evidence indicates that the ancestor of *C. elegans* was obligately outcrossing with separate male and female individuals (Kiontke & Fitch, 2005). Thus, *C. remanei* provides a useful proxy for the ancestral state of *C. elegans*, and comparisons between the two species provide a way of detecting changes in the evolutionary pressures experienced by *C. elegans* upon adopting a self-fertilizing breeding system. Analyses of sequence polymorphism show that *C. elegans* has much lower levels of genetic variation than *C. remanei* (Graustein *et al.*, 2002; Jovelin *et al.*,

2003; Haag & Ackerman, 2005; Cutter, 2006, 2008; Cutter *et al.*, 2006). *C. remanei* also suffers from much stronger inbreeding depression than *C. elegans* (Dolgin *et al.*, 2007), and the two species display very different mating behaviours (Chasnov *et al.*, 2007; Garcia *et al.*, 2007). Here, we show that natural populations of *C. elegans* and *C. remanei* also show markedly different transposon frequency distributions. The greater proportion of polymorphic insertions in *C. remanei* segregating at lower population frequencies compared with *C. elegans* is consistent with an important role of breeding systems contributing to the control of transposon dynamics in natural populations, and with other comparative studies of TEs in the plant genera *Arabidopsis* (Wright *et al.*, 2001) and *Solanum* (Tam *et al.*, 2007).

The estimates of N_{es} reveal distinctly different intensities of natural selection against transposons in *C. elegans* and *C. remanei*. We found signs of purifying selection against element mTcrl insertions in *C. remanei*, as indicated by negative N_{es} values, but no such evidence for Tc1 elements in *C. elegans*. It should be noted, however, that the elements examined in each species are not strictly homologous, which could potentially influence the contrast of TE polymorphism patterns between the two species. Since the unique internal sequence of mTcrl elements probably does not contain an open reading frame, these transposons are probably not autonomous and might experience different dynamics from the autonomous Tc1 elements. However, one would expect non-autonomous elements to be less mobile, and thus more likely to be fixed or at high frequencies (Bartolomé & Maside, 2004), whereas we see the opposite result. This potential innate difference between the elements would make our species contrast conservative. Alternatively, selection for self-regulation in selfing lineages could drive lower transposition rates in *C. elegans* (Charlesworth & Langley, 1986), or the *effective* transposition rate might be lower in *C. elegans* if transposons cannot easily invade and spread into different genetic backgrounds with self-fertilization. If transposition rates are lower for Tc1 elements than for mTcrl elements, fewer elements would be of recent origin in *C. elegans*, and population frequencies could be skewed upwards. In *C. remanei*, on the other hand, if selection effectively removes most elements, we might expect only to find either newly transposed elements at low frequencies or to observe old insertions that achieved high frequencies by drift. This would be consistent with the somewhat bimodal distribution for mTcrl elements seen in Fig. 3.

At first sight, our analysis of the intensity of selection suggests neutrality or positive selection for Tc1 elements in *C. elegans*. Although TEs in general can sometimes be beneficial (Kidwell & Lisch, 2001;

Schlenke & Begun, 2004), this is unlikely to be generally true for Tc1 elements, which are known to cause strong deleterious mutational effects (Plasterk & van Luenen, 1997; Bégin & Schoen, 2006, 2007). What might bias $N_{e,s}$ estimates for *C. elegans*? One possibility is violation of the assumption of copy number equilibrium required by the maximum likelihood method used for calculating the intensity of selection, as estimates of nucleotide diversity are lower than equilibrium predictions. However, the low estimates of genetic diversity between insertions in N2 could potentially result from gene conversion among Tc1 elements, thus biasing the molecular diversity estimates. The observation that Tc1 elements can acquire the sequence of other Tc1 elements elsewhere in the genome suggests that there might be continuous exchange of sequence information between individual insertions (Fischer *et al.*, 2003). But if between elements gene conversion is unbiased, theory shows that this should not affect diversity measures (Ohta, 1985; Slatkin, 1985). We cannot exclude the possibility that gene conversion is biased, which could reduce diversity. However, there is no reason why this should cause the excess of low-frequency variants that we detected as a negative Tajima's D , because gene conversion would introduce variants randomly across the genealogy (Marais, 2003).

The greater proportion of sites in linkage disequilibrium in *C. elegans* also suggests that the assumption of independence between insertions might be violated; although it is interesting to note that the levels of linkage disequilibrium reported here between TEs are much lower than the estimates of linkage disequilibrium in *C. elegans* from sequence data (Cutter, 2006, 2008; Cutter *et al.*, 2006) or other molecular markers (Barrière & Félix, 2005, 2007; Haber *et al.*, 2005). This could reflect a difference in timescale between transposons and other mutational processes. If transposition rates greatly exceed mutation rates, then transposon insertions will tend to be more recent, making them more likely to be independent of each other. This idea is supported by the significantly negative value for Tajima's D , which is also consistent with a recent burst of transposition (Sánchez-García *et al.*, 2005). However, this would then imply that strains lacking germline Tc1 activity only recently acquired suppressors of transposition (Collins *et al.*, 1987; Babity *et al.*, 1990; Mori *et al.*, 1990), and the presence of numerous fixed and high-frequency transposons indicates that many insertions preceded strain divergence. If we compare silent-site diversity among Tc1 elements from the N2 strain between fixed and near-fixed ($\geq 95\%$ population frequency) insertions with polymorphic insertions, we observe a nearly two-fold difference in diversity levels (fixed/near-fixed: $\pi_{si} = 0.31\%$ and $\theta_{si} = 0.73\%$; polymorphic: $\pi_{si} = 0.17\%$ and $\theta_{si} = 0.41\%$), further

suggesting that the high-frequency elements are more ancient. One explanation for this pattern is that there was a period of transpositional activity in the past, with the resulting insertions drifting to high frequencies or fixation, and the lower frequency TEs seen in our study represent a more recent period of activity. This resembles the pattern seen in *Drosophila melanogaster* (Bartolomé & Maside, 2004).

One scenario to explain the high number of fixed or near-fixed insertions across all chromosomes in *C. elegans* is that a nearly genome-wide selective sweep of an N2-like genotype with low levels of excision occurred in a number of strains. Such an event, however, would be unlikely to account for around half the insertions that show intermediate frequencies. Further, the complete lack of low-frequency ($< 20\%$) insertions in *C. elegans*, compared with around half the elements with low frequencies in *C. remanei* (see Fig. 3), suggests that the observed pattern is more likely to reflect differences in selection pressures between the species. Therefore, we caution against drawing strong conclusions from the positive estimates of selection for *C. elegans* from this analysis, because of the likelihood that population demographic processes might skew the transposon frequency spectrum. Rather, we argue that segregating Tc1 elements are probably selectively neutral, as seen in the analysis of the Hermanville population and some of the scattered samples.

Two main factors may be involved in causing the observed differences in transposon profiles. First, the skewed distribution in *C. elegans* towards high-frequency elements suggests a reduction in the efficacy of purifying selection in selfing lineages, due to a smaller effective population size (Wright & Schoen, 1999; Morgan, 2001). New transposons may persist longer in a polymorphic state in *C. remanei* and selection should be more efficient at eliminating them, because its effective population size is estimated to be two orders of magnitude larger than that of *C. elegans* (Cutter, 2006, 2008; Cutter *et al.*, 2006). Assuming that transposons have similar selective effects in both species, such large population size differences could explain the different $N_{e,s}$ estimates. But transposons could experience different selective regimes due to differences in levels of homozygosity between the breeding systems. When homozygous and heterozygous insertions have distinct fitness effects, changes in selfing rate can have dramatic effects on TE dynamics (Wright & Schoen, 1999; Morgan, 2001). Under the ectopic exchange model, selection against homozygous insertions is expected to be weak or null, whereas under the deleterious insertion model, selection will be strongest against homozygous insertions (see the review by Nuzhdin, 1999). Heterozygosity levels in *C. elegans* are typically very low (Barrière & Félix, 2005, 2007; but see Sivasundar & Hey, 2005),

suggesting that most insertions will be in a homozygous state, which would reduce the opportunity for ectopic pairing (Montgomery *et al.*, 1991). Therefore, a second explanation for the high population frequencies of elements in *C. elegans* is that selection is weaker against the greater proportion of homozygous insertions, as predicted by the ectopic exchange model.

Previous studies of polymorphic Tc1 elements in *C. elegans* have compared the canonical N2 strains with only a limited number of natural isolates (Emmons *et al.*, 1983; Liao *et al.*, 1983; Eide & Anderson, 1985; Harris & Rose, 1989; Egilmez *et al.*, 1995; Hodgkin & Doniach, 1997). Here, we studied *C. elegans* strains representing a complete global sampling covering all six major continents where *C. elegans* has been found, as well as a number of strains derived from a single geographic locality. Nevertheless, we observe many similarities in our dataset to those observed earlier for only a handful of strains. Egilmez *et al.* (1995) assessed Tc1-site occupancy among five strains, including N2, and found that 20 of the 32 Tc1 insertions in N2 were common to all five strains. With our larger sample of 40 strains, we also observed high numbers of fixed insertions (9 of 32). Following the observation that the Bergerac strain had active germline transposition and many times more Tc1 elements than N2 (Emmons *et al.*, 1983; Liao *et al.*, 1983), much of the focus on describing natural variation in Tc1 elements has been on characterizing strains as either ‘low-copy’ (~30 elements) or ‘high-copy’ (>300 elements) strains. We have not quantified copy numbers in the present study, and not all the strains used have been characterized previously for transposon abundance, but it is interesting to note that the two high-copy strains identified previously, RW7000/Bergerac and TR403 (Egilmez *et al.*, 1995; Hodgkin & Doniach, 1997), were also among the strains containing the highest fraction of insertions present (see Fig. 2A). This confirms previous findings that nearly all of the Tc1 elements in N2 are in the same location in the high-copy Bergerac strain (Harris & Rose, 1989).

An outstanding question that cannot be answered by our method is how the total numbers of transposons in natural isolates compare within and between the two species. In *C. elegans*, Tc1 copy number ranges an order of magnitude between wild strains (Egilmez *et al.*, 1995; Hodgkin & Doniach, 1997). In a single *C. remanei* genome (PB4641), we identified 81 mTc1 elements by bioinformatics approaches, although this is likely to underestimate the true copy number, and we do not know how copy numbers vary among strains. Furthermore, it is unclear how biologically relevant it is to compare the total genomic abundance of Tc1 and mTc1 elements if they are not homologous. Previous studies in related self- and cross-pollinating plant species similarly found that

outcrossing species consistently showed lower population frequencies of elements (Wright *et al.*, 2001; Tam *et al.*, 2007), but not of element copy number. Whereas self-fertilizing *Arabidopsis* had slightly higher copy numbers of DNA transposons than outcrossing species (Wright *et al.*, 2001), no relationship was found between retrotransposon number and breeding system among related tomato species (Tam *et al.*, 2007). Additional comparisons of TEs in other closely related species or using other TE families should help illuminate the effect of breeding system on total abundance. For *Caenorhabditis*, more work is needed to characterize and quantify transposons in *C. remanei* in particular.

The results presented here support a role of breeding systems in driving TE dynamics. Two independent derivations of self-fertilization in *Caenorhabditis* (Kiontke & Fitch, 2005) among the six species with sequenced genomes will provide a useful platform for testing the generality of breeding system as an important factor in TE evolution. Future studies will help to further tease apart the selection pressures imposed by TEs, and the impact of selfing rates on TE dynamics.

We are grateful to D. Petrov for sharing Mathematica files for the selection analysis, and to A. Betancourt and S. Wright for helpful comments on the manuscript. Worm strains were generously provided by S. Baird, M.-A. Félix and the *Caenorhabditis* Genetics Center. This work was funded by a Scottish International Education Trust travel grant, and by postgraduate scholarships from the Natural Sciences and Engineering Research Council of Canada (NSERC) and the University of Edinburgh School of Biological Sciences to E.S.D., and by an NSERC Discovery Grant to A.D.C. B.C. was supported by the Royal Society (UK).

References

- Abad, P., Quiles, C., Tares, S., Piotte, C., Castagnone-Sereno, P., Abadon, M. & Dalmasso, A. (1991). Sequences homologous to Tc(s) transposable elements of *Caenorhabditis elegans* are widely distributed in the phylum Nematoda. *Journal of Molecular Evolution* **33**, 251–258.
- Babity, J. M., Starr, T. V. B. & Rose, A. M. (1990). Tc1 transposition and mutator activity in a Bristol strain of *Caenorhabditis elegans*. *Molecular and General Genetics* **222**, 606–611.
- Barrière, A. & Félix, M.-A. (2005). High local genetic diversity and low outcrossing rate in *Caenorhabditis elegans* natural populations. *Current Biology* **15**, 1176–1184.
- Barrière, A. & Félix, M.-A. (2007). Temporal dynamics and linkage disequilibrium in natural *Caenorhabditis elegans* populations. *Genetics* **176**, 999–1011.
- Bartolomé, C. & Maside, X. (2004). The lack of recombination drives the fixation of transposable elements on the fourth chromosome of *Drosophila melanogaster*. *Genetical Research* **83**, 91–100.
- Bartolomé, C., Maside, X. & Charlesworth, B. (2002). On the abundance and distribution of transposable elements

- in the genome of *Drosophila melanogaster*. *Molecular Biology and Evolution* **19**, 926–937.
- Batzler, M. A. & Deininger, P. L. (2002). *Alu* repeats and human genomic diversity. *Nature Reviews Genetics* **3**, 370–379.
- Bennett, E. A., Coleman, L. E., Tsui, C., Pittard, W. S. & Devine, S. E. (2004). Natural genetic variation caused by transposable elements in humans. *Genetics* **168**, 933–951.
- Bessereau, J.-L. (2006). Transposons in *C. elegans*. In *Wormbook* (ed. The *C. elegans* Research Community), <http://www.wormbook.org>, doi/10.1895/wormbook.1.70.1.
- Boissinot, S., Entezam, A. & Furano, A. V. (2001). Selection against deleterious LINE-1-containing loci in the human lineage. *Molecular Biology and Evolution* **18**, 926–935.
- Boulesteix, M., Simard, F., Antonio-Nkondjio, C., Awono-Ambene, H. P., Fontenille, D. & Biémont, C. (2007). Insertion polymorphism of transposable elements and population structure of *Anopheles gambiae* M and S molecular forms in Cameroon. *Molecular Ecology* **16**, 441–452.
- Brookfield, J. F. (1986). A model for DNA sequence evolution within transposable element families. *Genetics* **112**, 393–407.
- Brookfield, J. F. Y. & Badge, R. M. (1997). Population genetic models of transposable elements. *Genetica* **100**, 281–294.
- Bégin, M. & Schoen, D. J. (2006). Low impact of germline transposition on the rate of mildly deleterious mutation in *Caenorhabditis elegans*. *Genetics* **174**, 2129–2136.
- Bégin, M. & Schoen, D. J. (2007). Transposable elements, mutational correlations, and population divergence in *Caenorhabditis elegans*. *Evolution* **61**, 1062–1070.
- Charlesworth, B. & Charlesworth, D. (1983). The population dynamics of transposable elements. *Genetical Research* **42**, 1–27.
- Charlesworth, D. & Charlesworth, B. (1995). Transposable elements in inbreeding and outbreeding populations. *Genetics* **140**, 415–417.
- Charlesworth, B. & Langley, C. H. (1986). The evolution of self-regulated transposition of transposable elements. *Genetics* **112**, 359–383.
- Charlesworth, B. & Langley, C. H. (1989). The population genetics of *Drosophila* transposable elements. *Annual Reviews of Genetics* **23**, 251–287.
- Chasnov, J. R., So, W. K., Chan, C. M. & Chow, K. L. (2007). The species, sex, and stage specificity of a *Caenorhabditis* sex pheromone. *Proceedings of the National Academy of Sciences of the USA* **104**, 6730–6735.
- Collins, J., Saari, B. & Anderson, P. (1987). Activation of a transposable element in the germ line but not the soma of *Caenorhabditis elegans*. *Nature* **328**, 726–728.
- Cutter, A. D. (2006). Nucleotide polymorphism and linkage disequilibrium in wild populations of the partial selfer *Caenorhabditis elegans*. *Genetics* **172**, 171–184.
- Cutter, A. D. (2008). Multilocus patterns of polymorphism and selection across the X-chromosome of *Caenorhabditis remanei*. *Genetics* **178**, 1661–1672.
- Cutter, A. D. & Payseur, B. A. (2003). Selection at linked sites in the partial selfer *Caenorhabditis elegans*. *Molecular Biology and Evolution* **20**, 665–673.
- Cutter, A. D., Baird, S. E. & Charlesworth, D. (2006). High nucleotide polymorphism and rapid decay of linkage disequilibrium in wild populations of *Caenorhabditis remanei*. *Genetics* **174**, 901–913.
- Denver, D. R., Morris, K. & Thomas, W. K. (2003). Phylogenetics in *Caenorhabditis elegans*: an analysis of divergence and outcrossing. *Molecular Biology and Evolution* **20**, 393–400.
- Dolgin, E. S., Charlesworth, B., Baird, S. E. & Cutter, A. D. (2007). Inbreeding and outbreeding depression in *Caenorhabditis* nematodes. *Evolution* **61**, 1339–1352.
- Dolgin, E. S., Félix, M.-A. & Cutter, A. D. (2008). Hakuna Nematoda: genetic and phenotypic diversity in African isolates of *Caenorhabditis elegans* and *C. briggsae*. *Heredity* **100**, 304–315.
- Doolittle, W. F. & Sapienza, C. (1980). Selfish genes, the phenotype paradigm and genome evolution. *Nature* **284**, 601–603.
- Duret, L., Marais, G. & Biémont, C. (2000). Transposons but not retrotransposons are located preferentially in regions of high recombination rate in *Caenorhabditis elegans*. *Genetics* **156**, 1661–1669.
- Duvernell, D. D. & Turner, B. J. (1999). Variation and divergence of death valley pupfish populations at retrotransposon-defined loci. *Molecular Biology and Evolution* **16**, 363–371.
- Egilmez, N. K., Ebert, R. H. II & Shmookler Reis, R. J. (1995). Strain evolution in *Caenorhabditis elegans*: transposable elements are markers of interstrain evolutionary history. *Journal of Molecular Evolution* **40**, 372–381.
- Eide, D. & Anderson, P. (1985). Transposition of Tc1 in the nematode *Caenorhabditis elegans*. *Proceedings of the National Academy of Sciences of the USA* **82**, 1756–1760.
- Emmons, S. W. & Yesner, L. (1984). High-frequency excision of transposable element Tc1 in the nematode *Caenorhabditis elegans* is limited to somatic cells. *Cell* **36**, 599–605.
- Emmons, S. W., Yesner, L., Ruan, K. S. & Katzenberg, D. (1983). Evidence for a transposon in *Caenorhabditis elegans*. *Cell* **32**, 55–65.
- Finnegan, D. J. (1992) Transposable elements. In *The Genome of Drosophila melanogaster* (ed. D. L. Lindsley & G. Zimm), pp. 1096–1107. New York: Academic Press.
- Fischer, S. E. J., Wienholds, E. & Plasterk, R. H. A. (2003). Continuous exchange of sequence information between dispersed Tc1 transposons in the *Caenorhabditis elegans* genome. *Genetics* **164**, 127–134.
- Franchini, L. F., Ganko, E. W. & McDonald, J. F. (2004). Retrotransposon-gene associations are widespread among *D. melanogaster* populations. *Molecular Biology and Evolution* **21**, 1323–1331.
- Garcia, L. R., LeBoeuf, B. & Koo, P. (2007). Diversity in mating behavior of hermaphroditic and male–female *Caenorhabditis* nematodes. *Genetics* **175**, 1761–1771.
- Graustein, A., Gaspar, J. M., Walters, J. R. & Palopoli, M. F. (2002). Levels of DNA polymorphism vary with mating system in the nematode genus *Caenorhabditis*. *Genetics* **161**, 99–107.
- Haag, E. S. & Ackerman, A. D. (2005). Intraspecific variation in *fem-3* and *tra-2*, two rapidly coevolving nematode sex-determining genes. *Gene* **349**, 35–42.
- Haber, M., Schüngel, M., Putz, A., Müller, S., Hasert, B. & Schulenburg, H. (2005). Evolutionary history of *Caenorhabditis elegans* inferred from microsatellites: evidence for spatial and temporal genetic differentiation and the occurrence of outbreeding. *Molecular Biology and Evolution* **22**, 160–173.
- Harris, L. J. & Rose, A. M. (1989). Structural analysis of Tc1 elements in *Caenorhabditis elegans* var. Bristol (strain N2). *Plasmid* **22**, 10–21.
- Harris, L. J., Prasad, S. & Rose, A. M. (1990). Isolation and sequence analysis of *Caenorhabditis briggsae* repetitive

- elements related to the *Caenorhabditis elegans* transposon Tc1. *Journal of Molecular Evolution* **30**, 359–369.
- Hickey, D. A. (1982). Selfish DNA: a sexually-transmitted nuclear parasite. *Genetics* **101**, 519–531.
- Hodgkin, J. & Doniach, T. (1997). Natural variation and copulatory plug formation in *Caenorhabditis elegans*. *Genetics* **146**, 149–164.
- Hoekstra, R., Otsen, M., Lenstra, J. A. & Roos, M. H. (1999). Characterisation of a polymorphic Tc1-like transposable element of the parasitic nematode *Haemonchus contortus*. *Molecular and Biochemical Parasitology* **102**, 157–166.
- Jacobson, J. W., Medhora, M. M. & Hartl, D. L. (1986). Molecular structure of a somatically unstable transposable element in *Drosophila*. *Proceedings of the National Academy of Sciences of the USA* **83**, 8684–8688.
- Jovelin, R., Ajie, B. C. & Phillips, P. C. (2003). Molecular evolution and quantitative variation for chemosensory behaviour in the nematode genus *Caenorhabditis*. *Molecular Ecology* **12**, 1325–1337.
- Ketting, R. F., Haverkamp, T. H., van Luenen, H. G. A. M. & Plasterk, R. H. A. (1999). *mut-7* of *C. elegans*, required for transposon silencing and RNA interference, is a homolog of Werner syndrome helicase and RNaseD. *Cell* **99**, 133–141.
- Kidwell, M. G. & Lisch, D. R. (2001). Transposable elements, parasitic DNA, and genome evolution. *Evolution* **55**, 1–24.
- Kiontke, K. & Fitch, D. H. A. (2005). The phylogenetic relationships of *Caenorhabditis* and other rhabditids. In *Wormbook* (ed. The *C. elegans* Research Community), <http://www.wormbook.org>, doi/10.1895/wormbook.1.11.1.
- Lander, E. S., Linton, L. M., Birren, B., Nusbaum, C., Zody, M. C., Baldwin, J., Devon, K., Dewar, K., Doyle, M., Fitzhugh, W., et al. (2001). Initial sequencing and analysis of the human genome. *Nature* **409**, 860–921.
- Liao, L. W., Rosenzweig, B. & Hirsh, D. (1983). Analysis of a transposable element in *Caenorhabditis elegans*. *Proceedings of the National Academy of Sciences of the USA* **80**, 3585–3589.
- Marais, G. (2003). Biased gene conversion: implications for genome and sex evolution. *Trends in Genetics* **19**, 330–338.
- Matsen, F. A. & Wakeley, J. (2006). Convergence to the island-model coalescent process in populations with restricted migration. *Genetics* **172**, 701–708.
- Medstrand, P., van de Lagemaat, L. N. & Mager, D. L. (2002). Retroelement distributions in the human genome: variations associated with age and proximity to genes. *Genome Research* **12**, 1483–1495.
- Moerman, D. G. & Waterston, R. H. (1989). Mobile elements in *Caenorhabditis elegans* and other nematodes. In *Mobile DNA* (ed. D. E. Berg & M. H. Howe), pp. 537–556. Washington, DC: American Society for Microbiology.
- Montgomery, E. A., Huang, S. M., Langley, C. H. & Judd, B. H. (1991). Chromosome rearrangement by ectopic recombination in *Drosophila melanogaster*: genome structure and evolution. *Genetics* **129**, 1085–1098.
- Morgan, M. T. (2001). Transposable element number in mixed mating populations. *Genetical Research* **77**, 261–275.
- Mori, I., Moerman, D. G. & Waterston, R. H. (1990). Interstrain crosses enhance excision of Tc1 transposable elements in *Caenorhabditis elegans*. *Molecular and General Genetics* **220**, 251–255.
- Neafsey, D. E., Blumenstiel, J. P. & Hartl, D. L. (2004). Different regulatory mechanisms underlie similar transposable element profiles in pufferfish and fruitflies. *Molecular Biology and Evolution* **21**, 2310–2318.
- Nuzhdin, S. V. (1999). Sure facts, speculations, and open questions about evolution of transposable elements. *Genetica* **107**, 129–137.
- Ohta, T. (1985). A model of duplicative transposition and gene conversion for repetitive DNA families. *Genetics* **110**, 513–524.
- Orgel, L. E. & Crick, F. H. (1980). Selfish DNA: the ultimate parasite. *Nature* **284**, 604–607.
- O'Brochta, D. A., Subramanian, R. A., Orsetti, J., Peckham, E., Nolan, N., Arensburger, P., Atkinson, P. W. & Charwood, D. J. (2006). *hAT* element population genetics in *Anopheles gambiae* s.l. in Mozambique. *Genetica* **127**, 185–198.
- Petrov, D., Aminetzrach, Y. T., Davis, J. C., Bensasson, D. & Hirsh, A. E. (2003). Size matters: non-LTR retrotransposable elements and ectopic recombination in *Drosophila*. *Molecular Biology and Evolution* **20**, 880–892.
- Plasterk, R. H. A., Izsvák, Z. & Ivics, Z. (1999). Resident aliens: the Tc1/*mariner* superfamily of transposable elements. *Trends in Genetics* **15**, 326–332.
- Plasterk, R. H. A. & van Luenen, H. G. A. M. (1997). Transposons. In *C. elegans II* (ed. D. L. Riddle, T. Blumenthal, B. J. Meyer & J. R. Priess), pp. 97–116. Cold Spring Harbor, NY: Cold Spring Harbor Laboratory Press.
- Rezsohazy, R., van Luenen, H. G. A. M., Durbin, R. M. & Plasterk, R. H. A. (1997). Tc7, a Tc1-hitch hiking transposon in *Caenorhabditis elegans*. *Nucleic Acids Research* **25**, 4048–4054.
- Rizzon, C., Marais, G., Gouy, M. & Biémont, C. (2002). Recombination rate and the distribution of transposable elements in the *Drosophila melanogaster* genome. *Genome Research* **12**, 400–407.
- Rizzon, C., Martin, E., Marais, G., Duret, L., Segalat, L. & Biémont, C. (2003). Patterns of selection against transposons inferred from the distribution of Tc1, Tc3 and Tc5 insertions in the *mut-7* line of the nematode *Caenorhabditis elegans*. *Genetics* **165**, 1127–1135.
- Rozas, J., Sanchez-DelBarrio, J. C., Messeguer, X. & Rozas, R. (2003). DnaSP, DNA polymorphism analyses by the coalescent and other methods. *Bioinformatics* **19**, 2496–2497.
- Sánchez-García, A., Maside, X. & Charlesworth, B. (2005). High rate of horizontal transfer of transposable elements in *Drosophila*. *Trends in Genetics* **21**, 200–203.
- Schlenke, T. A. & Begun, D. J. (2004). Strong selective sweep associated with a transposon insertion in *Drosophila simulans*. *Proceedings of the National Academy of Sciences of the USA* **101**, 1626–1631.
- Sivasundar, A. & Hey, J. (2005). Sampling from natural populations with RNAi reveals high outcrossing and population structure in *Caenorhabditis elegans*. *Current Biology* **15**, 1598–1602.
- Slatkin, M. (1985). Genetic differentiation of transposable elements under mutation and unbiased gene conversion. *Genetics* **110**, 145–158.
- Takasaki, N., Amaki, T. Y., Hamada, M., Park, L. & Okada, N. (1997). The salmon *SmaI* family of short interspersed repetitive elements (SINES): interspecific and intraspecific variation of the insertion of SINES in the genomes of chum and pink salmon. *Genetics* **146**, 369–380.

- Tam, S. M., Causse, M., Garchery, C., Burck, H., Mhiri, C., & Granbastien, M.-A. (2007). The distribution of *copia* -type retrotransposons and the evolutionary history of tomato and related wild species. *Journal of Evolutionary Biology* **20**, 1056–1072.
- van Luenen, H. G. A. M., Colloms, S. D. & Plasterk, R. H. A. (1994). The mechanism of transposition of Tc3 in *C. elegans*. *Cell* **79**, 293–301.
- Vos, J. C. & Plasterk, R. H. A. (1994). Tc1 transposase of *Caenorhabditis elegans* is an endonuclease with a bipartite DNA binding domain. *EMBO Journal* **13**, 6125–6132.
- Wakeley, J. (2003). Polymorphism and divergence for island-model species. *Genetics* **163**, 411–420.
- Weir, B. S. & Hill, W. G. (1980). Effect of mating structure on variation in linkage disequilibrium. *Genetics* **95**, 477–488.
- Wright, S. I. & Schoen, D. J. (1999). Transposon dynamics and the breeding system. *Genetica* **107**, 139–148.
- Wright, S. I., Hien Le, Q., Schoen, D. J. & Bureau, T. E. (2001). Population dynamics of an *Ac*-like transposable element in self- and cross-pollinating *Arabidopsis*. *Genetics* **158**, 1279–1288.
- Zampicini, G., Blinov, A., Cervella, P., Guryev, V. & Sella, G. (2004). Insertional polymorphism of a non-LTR mobile element (NLRCth1) in European populations of *Chironomus riparius* (Diptera, Chironomidae) as detected by transposon insertion display. *Genome* **47**, 1154–1163.